

## References

- Batteau, D. W. (1967). The role of the pinna in human localization. *Proc. Roy. Soc. (London B)* **158**: 158–180.
- de Boer, E. (1956). Pitch of inharmonic signals. *Nature (London)* **178**: 535–536.
- Carr, C. E. and Konishi, M. (1988). Axonal delay lines for time measurement in the owl's brainstem. *Proc. Natl. Acad. Sci.* **85**: 8311–8315.
- Dallos, P. (1985). Response characteristics of mammalian cochlear hair cells. *J. Neurosci.* **5**: 1591–1608.
- Deutsch, D. (1981). The octave illusion and auditory perceptual integration. In Tobias, J. V. (ed). *Hearing Research and Theory*, Vol. I. New York: Academic Press, pp. 99–142.
- Durlach, N. I. (1972). Binaural signal detection: Equalization and cancellation theory. In Tobias, J. V. (ed). *Foundations of Modern Auditory Theory*, Vol. II. New York: Academic Press, pp. 371–462.
- Evans, E. F. (1982). Functional anatomy of the auditory system. In Barlow, H. B. and Mollon, J. D. (eds), *The Senses*. Cambridge, England: Cambridge University Press, p. 251.
- Fourcin, A.J. (1965). The pitch of noise with periodic spectral peaks. In *Proceedings 5th International Congress on Acoustics, Leige*, Vol. Ia, B 52.
- Fujita I., and Konishi, M. (unpublished). The role of GABAergic inhibition in coding of interaural time difference in the owl's auditory system.

- Geisler, C.D. and Greenberg, S. (1986). A two-stage nonlinear cochlear model posseses automatic gain control. *J. Acoust. Soc. Am.* **80**: 1359–1363.
- Goblick, T.J., Jr., and Pfeiffer, R. R. (1969) Time domain measurements of cochlear non-linearities using combination click stimuli. *J. Acoust. Soc. Am.* **46**: 924–938.
- Goldberg, J. M. and Brown, P. B. (1969). Response of binaural neurons to dog superior olivary complex to dichotic tonal stimuli: Some physiological mechanisms of sound localization. *J. Neurophysiol.* **32**: 613–636.
- Goldstein, J. L. (1973). An optimum processor theory for the central formation of the pitch of complex tones. *J. Acoust. Soc. Am.* **54**: 1496–1516.
- Greenberg, S. (1988). The ear as a speech analyzer. *J. Phonetics* **16**: 139–149.
- Helmholtz, H. L. F. (1895). *Sensations of Tone*. Ellis, A. J. (Trans.) New York: Longmans Green, pp. 49–65.
- Jeffress, L. A. (1948). A place theory of sound localization. *J. Comp. Physiol. Pyschol.* **41**: 35–39.
- Jeffress, L. A. (1972). Binaural signal detection: Equalization and cancellation theory. In Tobias, J. V. (ed). *Foundations of Modern Auditory Theory*, Vol. II. New York: Academic Press, pp. 349–368.
- Kiang, N. Y.-s, Watenabe, T., Thomas, E.C., and Clark, L.F. (1965). *Discharge Patterns of Single Fibers in the Cat's Auditory Nerve*. Cambridge, MA: MIT Press.
- Kiang, N. Y.-s (1980). Processing of speech by the auditory nervous system. *J. Acoust. Soc. Am.* **68**: 830–835.

Kim, D. O. (1984). Functional roles of the inner- and outer-haircell subsystems in the cochlea and brainstem. In Berlin, C. I. (ed), *Hearing Science*. San Diego, CA: College-Hill Press, pp. 241–262.

Knudsen, E. I. and Konishi, M. (1978). A neural map of auditory space in the owl. *Science* **200**: 795–797.

Knudsen, E. I., Blasdel, G. G., and Konishi, M. (1979). Sound localization by the barn owl measured with the search coil technique. *J. Comp. Physiol.* **133**: 1–11.

Knudsen, E. I. and Konishi, M. (1979). Mechanisms of sound localization in the barn owl (*Tyto alba*). *J. Comp. Physiol.* **133**: 13–21.

Knudsen, E. I. and Konishi, M. (1980). Monaural occlusion shifts receptive field locations of auditory midbrain units in the owl. *J. Neurophysiol* **44**: 687–695.

Konishi, M. (1986). Centrally synthesized maps of sensory space. *TINS* **4**: 163–168.

Lazzaro, J. P., Ryckebusch, S., Mahowald, M. A., and Mead, C. (1988). Winner-take-all networks of  $O(n)$  complexity. In Touretzky, D. (ed), *Advances in Neural Information Processing Systems 1*. San Mateo, CA: Morgan Kaufmann Publishers, pp. 703-711.

Lazzaro, J. P. and Mead, C. (1989a). Silicon models of auditory localization. *Neural Computation* **1**: 41–70.

Lazzaro, J. P. and Mead, C. (1989b). Circuit models of sensory transduction in the cochlea. In Mead, C. and Ismail, M. (eds), *Analog VLSI Implementations of Neural Networks*. Norwell, MA: Kluwer Academic Publishers, pp. 85-101.

- Liberman, A. M. and Mattingly, I. G. (1989). A specialization for speech perception. *Science* **243**: 489–494.
- Licklider, J. C. R. (1951). A duplex theory of pitch perception. *Experientia* **7**: 128–134.
- Licklider, J. C. R. (1954). “Periodicity pitch” and “place pitch.” *J. Acoust. Soc. Am.* **26**: 945 (A).
- Licklider, J. C. R. (1959). Three auditory theories. In Koch, S. (ed), *Pyschology: A Study of a Science*, Vol 1. New York: McGraw-Hill, pp. 94–144.
- Lyon, R. F. (1984). Computational models of neural auditory processing. In *Proceedings, 1984 IEEE ICASSP*, San Diego, CA (Mar. 1984).
- Lyon, R. F. and Mead, C. (1988a). An analog electronic cochlea. *IEEE Trans. Acoust., Speech, Signal Processing* **36**: 1119–1134.
- Lyon, R. F. and Mead, C. (1988b). *Cochlear Hydrodynamics Demystified*. Caltech Computer Science Technical Report, Caltech-CS-TR-88-4, California Institute of Technology, Pasadena, CA, February 1988.
- Mahowald, M.A., and Delbruck, T.I. (1989). Cooperative stereo matching using static and dynamic image features. In Mead, C. and Ismail, M. (eds), *Analog VLSI Implementations of Neural Networks*. Norwell, MA: Kluwer Academic Publishers, pp. 181-206.
- Mead, C. (1989). *Analog VLSI and Neural Systems*. Reading, MA: Addison-Wesley.
- Manley, G. A., Koppl, C., and Konishi, M. (1988). A neural map of interaural intensity difference in the brain stem of the barn owl. *J. Neurosci.* **8**: 2665-2676.

- Moiseff, A. and Konishi, M. (1981). Neuronal and behavioral sensitivity to binaural time differences in the owl. *J. Neurosci.* **1**: 40–48.
- Moiseff, A. and Konishi, M. (1983). Binaural characteristics of units in the owl's brainstem auditory pathways: Precursors of restricted spatial fields. *J. Neurosci.* **3**: 2553–2562.
- Rhode, W. S. (1971). Observations of the vibration of the basilar membrane in squirrel monkeys using the Mossbauer technique. *J. Acoust. Soc. Am.* **49**: 1218–1231.
- Rhode, W. S., Geisler, C.D., and Kennedy, D.T. (1978). Auditory nerve fiber response to wide-band noise and tone combinations. *J. Neurophysiol.* **41**: 692–704.
- Robles, L., Rhode, W. S., and Geisler, C.D. (1976) Transient response of basilar membrane measured in squirrel monkeys using the Mossbauer effect. *J. Acoust. Soc. Am.* **59**: 926–939.
- Rose, J.E., Hind, J.E., Anderson, D. J., and Brugge, J. F. (1971). Some effects of stimulus intensity on response of auditory nerve fibers in the squirrel monkey. *J. Neurophysiol.* **34**: 685–699.
- Sachs, M. B. and Abbas, P. J. (1974) Rate versus level functions for auditory-nerve fibers in cats: Tone-burst stimuli. *J. Acoust. Soc. Am.* **56**: 1835–1847.
- Schouten, J. F. (1940). The residue, a new component in subjective sound analysis. *Proc. Kon. Acad. Wetensch. (Neth.)* **43**: 356–365.
- Suga, N. (1984). The extent to which biosonar information is represented in the bat auditory cortex. In Edelman, G. M., Gall, W. E., and Cowan, W. M. (eds), *Dynamic Aspects of Neocortical Function*. New York: Wiley, 1984.

- Sullivan, W. E. and Konishi, M. (1984). Segregation of stimulus phase and intensity coding in the cochlear nucleus of the barn owl. *J. Neurosci.* **7**: 1787–1799.
- Takahashi, T. T. and Konishi, M. (1988a). Projections of the cochlear nuclei and nucleus laminaris to the inferior colliculus of the barn owl. *J. Compar. Neurol.* **274**: 190–211.
- Takahashi, T. T. and Konishi, M. (1988b). Projections of the nucleus angularis and nucleus laminaris to the lateral lemniscal nuclear complex of the barn owl. *J. Compar. Neurol.* **274**: 221–238.
- Tobias, J. V. (1972). Curious binaural phenomena. In Tobias, J. V. (ed). *Foundations of Modern Auditory Theory*, Vol. II. New York: Academic Press, pp. 463–486.
- Wagner, H., and Konishi, M. (unpublished).
- Wightman, F. L. (1973). The pattern-transformation model of pitch. *J. Acoust. Soc. Am.* **54**: 407–416.
- Yin, T. C. T and Kuwada, S. (1984). Neuronal mechanisms of binaural interaction. In Edelman, G.M., Gall, W.E., and Cowan, W.M. (eds), *Dynamic Aspects of Neocortical Function*. New York: Wiley, 1984.